

Clustering around QSOs at $1.0 < z < 1.6$

Sebastián F. Sánchez^{1,2}, J.I. González-Serrano¹

¹*Instituto de Física de Cantabria-CSIC, Avd. De los Castros S/N, 39005-Santander, Spain*

²*Dept. de Física Moderna, Facultad de Ciencias, Univ. de Cantabria, Avd. De los Castros S/N, 39005-Santander*

Abstract. Deep B and R band images ($R < 24$ and $B < 25$) have been obtained of a field of $\sim 6 \times 6$ arcmin around seven radio-loud quasars with $1 < z < 1.6$. A statistically significant excess of *faint* ($22.5 < B \leq 23.5$ and $22 < R \leq 23.0$) galaxies was found, on scales of $r < 160''$ around the quasars ($> 3\sigma_{N_{\text{exp}}}$). The galaxies have magnitudes compatible with being at the redshift of the QSOs.

1 Introduction

Evidence has been accumulating over last decades that there is a link between the quasar activity and overdensities of galaxies around the host galaxy ([1], [2], [3], and references therein). Recent results, [3], have shown that there is an excess of *faint* galaxies around a sample of 31 radio-loud quasars at high- z ($z > 1$). The magnitudes and colors of the excess galaxies are consistent with a population of mostly early-type galaxies at the quasar redshift. The size of this overdensity has not been completely determined, being larger than $\theta \sim 100''$ (due to the size of the images, about $4' \times 4'$).

We report on the results obtained from deep ($B < \sim 25$ and $R < \sim 24$) CCD imaging survey of the fields around 7 B3VLA quasars, in the redshift range $1.0 < z < 1.6$. The images were obtained during the night of 1997 March 13 at the 2.2m Calar Alto telescope, with a final integration time of 1800 and 2400 seconds for the R and B band, respectively. The size of the images, $6' \times 6'$ have allowed us to sample the density of galaxies to scales down to $\sim 190''$.

2 Data processing and number counts

We searched for objects in the B and R band images using the SExtractor package. Our detection threshold was of 5 connected pixels above 2σ level per pixel, which guarantees a detection up to 4σ . Simulations have been performed to determine the limits of the validity of the method (detection+magnitude determination+star-galaxy classification) showing that it is valid down to $R \leq 23.5$ -23.8 and $B \leq 24.3$ -24.5.

The *field* number-counts and its variance have been obtained directly from the images, measuring the number of galaxies in an annulus from $r_{in}=170''$ to




Figure 1: Relative excess galaxies averaged over the seven quasar fields and both bands.

$r_{out}=185''$, and in a grid of 16 circular, not overlapping, areas of $r < 35''$ (centered more than $70''$ from the quasars). The measurements were repeated for different ranges of magnitudes, which define the number-counts distribution.

3 Results

Figure 1 shows the relative excess of galaxies averaged over the seven fields, at both B and R bands. There is a significant excess of galaxies around the quasars (5.52σ). The excess presents a tendency with the magnitude range, being significant only for the faintest magnitudes ($22 \leq R \leq 23$ and $22.5 \leq B \leq 23.5$), for any angular scale (down to $\theta \sim 160''$).

These results indicate most probably that the excess is not due to intervening galaxies, since there is *no* significant excess down to $R < 21$ and $B < 22.5$ mag. The magnitudes ($R \sim 22.5$ and $B \sim 23.0$), angular scale and number of galaxies of the excess are compatible with being clusters or groups of galaxies at the redshift of the quasars.

References

- [1] Yee H.K. & Green R.F., 1987, ApJ, 319, 28
- [2] Yee H.K. & Ellingson E., 1993, ApJ, 411, 43
- [3] Hall P.B. & Green R.F., 1998, ApJ, in press [SISSA pre-prints: astro-ph/9806151]